

What is claimed is:

1. A method of screening for compounds that inhibit melanogenesis, the method comprising: treating cells expressing a tyrosinase-encoding gene with a test compound, and
5 determining the cellular localization of tyrosinase in the presence of the test compound; wherein a change in the cellular localization of tyrosinase in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.
2. The method of claim 1, wherein the cells further express a P protein-encoding gene,
10 and wherein the change in the cellular localization of tyrosinase in the presence of the test compound as compared to in the absence of the test compound is dependent upon the expression of the P protein-encoding gene.
3. The method of claim 1 or 2, wherein the cellular localization of tyrosinase is determined
15 by assaying the amount of tyrosinase secreted by the cells in the presence of the compound, wherein an increase in the amount of tyrosinase secreted by the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.
4. The method of claim 1 or 2, wherein the cellular localization of tyrosinase is detected
by assaying for tyrosinase activity.
- 20 5. The method of claim 1 or 2, wherein the cellular localization of tyrosinase is detected by assaying for the presence of tyrosinase protein using immunological techniques.
6. The method of claim 1 or 2 further comprising the step of assaying the amount of tyrosinase associated in a high molecular weight complex in the presence of the test
25 compound, wherein a decrease in the amount of tyrosinase associated in a high molecular weight complex in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.
7. The method of claim 1 or 2 further comprising the step of assaying the amount of
30 TRP-1 or TRP-2 protein associated in a high molecular weight complex in the presence of the compound, wherein a decrease in the amount of TRP-1 or TRP-2 protein associated in a high molecular weight complex in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.

8. The method of claim 1 or 2 further comprising the step of assaying the number or size of melanosomes in the cells in the presence of the compound, wherein a decrease in the number or size of melanosomes in the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound inhibits melanogenesis.
9. The method of claim 1 or 2 further comprising the step of assaying the mass or length of tyrosinase in the cells in the presence of the compound, wherein a decrease in the mass or length of tyrosinase in the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.
10. The method of claim 1 or 2 further comprising the step of assaying for the levels and/or targeting of lysosomal hydrolases in the cells in the presence of the compound, wherein a decrease in accumulation of lysosomal hydrolases that are transported via the M6P/IGF-II receptor in the lysosome in the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that inhibits melanogenesis.
11. The method of claim 1 or 2, wherein the cells are grown in the presence of low tyrosine.
12. The method of claim 11 wherein the concentration of tyrosine is 0.01-0.03 mM.
13. The method of claim 1 wherein the cells are melanocytes.
14. The method of claim 1 wherein the cells are melanoma cells.
15. The method of claim 1 wherein the cells are derived from a mammal.
16. The method of claim 15 wherein the mammal is a human.
17. The method of claim 15 wherein the mammal is selected from the group consisting of mouse, hamster, and guinea pig.
18. A method of screening for compounds that increase melanogenesis comprising: treating cells expressing a tyrosinase-encoding gene with a test compound, and determining the amount of tyrosinase secreted by the cells in the presence of the test compound; wherein a decrease in the amount of tyrosinase secreted by the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound is a candidate for a compound that increases melanogenesis.

19. The method of claim 18 wherein the cells further express a P protein-encoding gene, and wherein the decrease in the amount of tyrosinase secreted by the cells in the presence of the test compound as compared to in the absence of the test compound is dependent upon the expression of the P protein-encoding gene.
- 5 20. The method of claim 18 or 19 further comprising determining a ratio of tyrosinase inside the cells to tyrosinase secreted by the cells, wherein an increase in the ratio in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound induces melanogenesis.
21. The method of claim 18 or 19, wherein the amount of tyrosinase is detected by
10 assaying for tyrosinase activity.
22. The method of claim 18 or 19, wherein the amount of tyrosinase is detected by assaying for the presence of tyrosinase protein using immunological techniques.
23. The method of claim 18 wherein the cells are melanocytes.
24. The method of claim 18 wherein the cells are melanoma cells.
- 15 25. The method of claim 23 or 24, wherein the cells are visually examined for an increase in melanin production.
26. The method of claim 23 or 24 wherein the cells do not express P protein, and wherein a decrease in the amount of tyrosinase secreted by the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test
20 compound is a candidate for a compound that mimics P protein function.
27. The method of claim 23 wherein the cells are mouse melan-p melanocytes.
28. The method of claim 18 wherein the cells are derived from a mammal.
29. The method of claim 28 wherein the mammal is a human.
30. The method of claim 28 wherein the mammal is selected from the group consisting of
25 mouse, hamster, and guinea pig.
31. The method of claim 26, wherein the cells are visually examined for an increase in melanin production.
32. A method of screening for compounds that affect the function of P protein, the method comprising: contacting a system with a test compound, the system comprising P protein and
30 tyrosinase; and identifying those test compounds that affect tyrosinase activity in the system in a P protein-dependent manner.

33. The method of claim 32 wherein the system is a cell that expresses a P protein-encoding gene and a tyrosinase-encoding gene.

34. The method of claim 33 wherein the cell is a cultured cell.

35. The method of claim 32 wherein compounds that decrease tyrosinase activity in the system are identified as compounds that inhibit the function of P protein.

36. The method of claim 35 further comprising the step of assaying for the targeting of lysosomal hydrolases in the cells in the presence of the compound, wherein a decrease in accumulation of lysosomal hydrolases that are transported via the M6P/IGF-II receptor in the lysosome in the cells in the presence of the test compound as compared to in the absence of the test compound indicates that the test compound inhibits the function of P protein.

37. The method of claim 32 wherein compounds that result in an increase in tyrosinase activity in the system are identified as compounds that increase the function of P protein.

38. The method of claim 32 wherein the P protein-encoding gene is derived from a mammal selected from the group consisting of human, hamster, guinea pig, and mouse.

39. The method of claim 32 wherein the tyrosinase-encoding gene is derived from a mammal selected from the group consisting of human, hamster, guinea pig, and mouse.

40. A method of screening for compounds that affect the function of P protein, the method comprising: using a primary screening method to preliminarily determine whether a test compound may affect P protein function; and using one or more secondary screening methods to determine whether the test compound affects P protein function.

41. The method of claim 40, wherein the primary screening method comprises at least one screening assay selected from the group consisting of assaying for secretion of tyrosinase and assaying for the missorting of at least one lysosomal hydrolase.

42. A method of screening for compounds that affect the function of P protein, the method comprising: modeling a compound that affects the function of the P protein; making chemical analogs of the compound; and assaying the chemical analogs for their effect on the function of P protein.

43. The method of claim 42 wherein the compound is imipramine.


44. A method of decreasing melanin production in a melanocyte, comprising contacting the melanocyte with an effective amount of a compound that effects an alteration in late endosomal/lysosomal trafficking in the melanocyte, wherein the alteration in late endosomal/lysosomal trafficking results in a decrease in melanin production in the melanocyte.

45. The method of claim 44, wherein the alteration in late endosomal/lysosomal trafficking is effected by contacting the melanocyte with a compound that is an antagonist of late endosomal/lysosomal trafficking.

46. The method of claim 44, wherein the alteration in late endosomal/lysosomal trafficking is an alteration in late endosomal/lysosomal cholesterol trafficking.

47. The method of claim 44, wherein the alteration in late endosomal/lysosomal trafficking is effected by contacting the melanocyte with a compound selected from the group consisting of

- (a) progesterone,
- (b) a hydrophobic amine,
- (c) sphingosine, and
- (d) a compound of the formula



wherein X is O or S;

R₁ is -C(O)(C₁-C₆)alkyl or -(CH₂)_n-O-(C₁-C₆)alkyl, or -(CH₂)_n-NR₇R₈ where n is 0-3, and where each of R₇ and R₈ are independently selected from H and (C₁-C₆)alkyl;

R₂ is H or (C₁-C₆)alkyl;

R₃ is H or (C₁-C₆)alkyl;

R₄ is -C(O)(C₁-C₆)alkyl;

R₅ is H or -(C₁-C₆)alkyl; or R₄ and R₅ together are =O; and

R₆ is H or -(C₁-C₆)alkyl or -(CH₂)_n-NR₉R₁₀ where each of R₉ and R₁₀ are independently selected from H and (C₁-C₆)alkyl.

48. The method of claim 47, wherein the compound is progesterone.


49. The method of claim 47, wherein the compound is a hydrophobic amine.

50. The method of claim 49, wherein the hydrophobic amine is selected from the group consisting of a phenothiazine, and a tricyclic antidepressant.





52. The method of claim 51, wherein the phenothiazine is selected from the group consisting of trifluoperazine, chlorpromazine, prochlorperazine, triflupromazine, promazine, thioridazine, mesoridazine, piperacetazine, perphenazine, fluphenazine, acetophenazine, and thiethylperazine.

54. The method of claim 53, wherein the tricyclic antidepressant is selected from the group consisting of imipramine, nortriptyline, protriptyline, trimipramine, and doxepin.


56. The method of claim 47, wherein the compound is selected from the group consisting of:



5



and



(VIII)

or a pharmaceutically acceptable salt or solvate thereof.


57. A method of reducing skin pigmentation, comprising contacting skin with a pharmaceutically effective amount of a compound that effects an alteration in late endosomal/lysosomal trafficking, wherein an alteration in late endosomal/lysosomal trafficking results in a reduction of skin pigmentation.

58. The method of claim 57, wherein an alteration in late endosomal/lysosomal trafficking is effected by contacting the skin with a compound that is an antagonist of late endosomal/lysosomal trafficking.

59. The method of claim 57, wherein the alteration in late endosomal/lysosomal trafficking is an alteration in late endosomal/lysosomal cholesterol trafficking.

60. The method of claim 57, wherein the alteration in late endosomal/lysosomal trafficking is effected by contacting the skin with a pharmaceutically effective amount of a compound selected from the group consisting of

- (a) progesterone,
- (b) a hydrophobic amine,
- (c) sphingosine, and
- (d) a compound of the formula



(I)

wherein X is O or S;

R₁ is -C(O)(C₁-C₆)alkyl or -(CH₂)_n-O-(C₁-C₆)alkyl, or -(CH₂)_n-NR₇R₈ where n is 0-3, and where each of R₇ and R₈ are independently selected from H and (C₁-C₆)alkyl,

R₂ is H or (C₁-C₆)alkyl;

R₃ is H or (C₁-C₆)alkyl;

R₄ is -C(O)(C₁-C₆)alkyl;

R₅ is H or -(C₁-C₆)alkyl; or R₄ and R₅ together are =O; and

5 R₆ is H or -(C₁-C₆)alkyl or -(CH₂)_n-NR₉R₁₀ where each of R₉ and R₁₀ are independently selected from H and (C₁-C₆)alkyl.

61. The method of claim 60, wherein the compound is progesterone.

62. The method of claim 60, wherein the compound is a hydrophobic amine.

63. The method of claim 62, wherein the hydrophobic amine is selected from the group
10 consisting of a phenothiazine, and a tricyclic antidepressant.

64. The method of claim 63, wherein the compound is a phenothiazine.


65. The method of claim 64, wherein the phenothiazine is selected from the group
consisting of trifluoperazine, chlorpromazine, prochlorperazine, triflupromazine, promazine,
thioridazine, mesoridazine, piperacetazine, perphenazine, fluphenazine, acetophenazine, and
15 thiethylperazine.

66. The method of claim 63, wherein the compound is a tricyclic antidepressant.





67. The method of claim 66, wherein the tricyclic antidepressant is selected from the group
consisting of imipramine, nortriptyline, protriptyline, trimipramine, and doxepin.

68. The method of claim 60, wherein the compound is sphingosine.


20 69. The method of claim 60, wherein the compound is selected from the group consisting
of:



5



103040-842350




5 70. A pharmaceutical composition for reducing skin pigmentation, comprising a skin pigmentation reducing effective amount of a compound that effects an alteration in late endosomal/lysosomal trafficking in a skin cell and a pharmaceutically acceptable carrier.

71. The composition of claim 70, wherein the alteration in late endosomal/lysosomal trafficking is effected by contacting the melanocyte with a compound that is an antagonist of late endosomal/lysosomal trafficking.

72. The composition of claim 70, wherein the alteration in late endosomal/lysosomal trafficking is an alteration in late endosomal/lysosomal cholesterol trafficking.

73. The composition of claim 70, wherein the compound that effects the alteration in late endosomal/lysosomal trafficking is selected from the group consisting of

- 15 (a) progesterone, 514/177
- (b) a hydrophobic amine,
- (c) sphingosine, and 514/669
- selected → (d) a compound of the formula



(I)

514/177, 482 etc
excludes progesterone

wherein X is O or S;

R₁ is -C(O)(C₁-C₆)alkyl or -(CH₂)_n-O-(C₁-C₆)alkyl, or -(CH₂)_n-NR₇R₈ where n is 0-3, and where each of R₇ and R₈ are independently selected from H and (C₁-C₆)alkyl;

5 R₂ is H or (C₁-C₆)alkyl;

R₃ is H or (C₁-C₆)alkyl;

R₄ is -C(O)(C₁-C₆)alkyl;

R₅ is H or -(C₁-C₆)alkyl; or R₄ and R₅ together are =O; and

10 R₆ is H or -(C₁-C₆)alkyl or -(CH₂)_n-NR₉R₁₀ where each of R₉ and R₁₀ are independently selected from H and (C₁-C₆)alkyl.

74. The composition of claim 73, wherein the compound is progesterone.

75. The composition of claim 73, wherein the compound is a hydrophobic amine.

76. The composition of claim 73, wherein the hydrophobic amine is selected from the group consisting of a phenothiazine, and a tricyclic antidepressant.

15 77. The composition of claim 76, wherein the compound is a phenothiazine.

78. The composition of claim 77, wherein the phenothiazine is selected from the group consisting of trifluoperazine, chlorpromazine, prochlorperazine, triflupromazine, promazine, thioridazine, mesoridazine, piperacetazine, perphenazine, fluphenazine, acetophenazine, and thiethylperazine.

20 79. The composition of claim 76, wherein the compound is a tricyclic antidepressant.

80. The composition of claim 79, wherein the tricyclic antidepressant is selected from the group consisting of imipramine, nortriptyline, protriptyline, trimipramine, and doxepin.

81. The composition of claim 73, wherein the compound is sphingosine.

25 82. The composition of claim 73, wherein the compound is selected from the group consisting of:

1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2

10





no antes bases ~
cl 13




Sub
con

T.03040-844266



and



5

83. A method of activating melanogenesis comprising contacting a melanocyte with diminished or absent P protein activity with a pharmaceutically effective amount of a compound that inhibits ATPase, whereby the inhibition of ATPase results in an activation of melanogenesis in the melanocyte with diminished or absent P protein activity.

10

84. The method according to claim 83, wherein the melanocyte with diminished or absent P protein activity is contacted with a pharmaceutically effective amount of a compound selected from the group consisting of bafilomycin, concanamycin, and derivatives thereof.

85. The method according to claim 83, wherein the melanocyte with diminished or absent P protein activity is contacted with a pharmaceutically effective amount of bafilomycin or a derivative thereof.

86. The method according to claim 83, wherein the melanocyte with diminished or absent
5 P protein activity is contacted with a pharmaceutically effective amount of concanamycin or a derivative thereof.

87. A method of treating tyrosinase-positive, oculocutaneous albinism in an individual in need thereof, the method comprising contacting skin of the individual with a pharmaceutically effective amount of a compound that inhibits ATPase, whereby the inhibition of ATPase results
10 in an activation of melanogenesis in the individual.

88. The method according to claim 87, wherein the skin of the subject is contacted with a pharmaceutically effective amount of a compound selected from the group consisting of bafilomycin or a derivative thereof and concanamycin or a derivative thereof.

89. The method according to claim 88, wherein the skin of the subject is contacted with a
15 pharmaceutically effective amount of bafilomycin or a derivative thereof.

90. The method according to claim 88, wherein the skin of the subject is contacted with a pharmaceutically effective amount of concanamycin or a derivative thereof.

91. A kit comprising a pharmaceutically effective amount of a compound that modulates melanogenesis by affecting P-protein function, inhibiting late endosomal/lysosomal trafficking, or inhibiting an ATPase.
20

92. The kit of claim 91, further comprising a set of written instructions describing how to use the compound to modulate skin pigmentation.

restrict
to extent they read on elected cpd.